

RESEARCH

Open Access

Effects of the standardized ileal digestible lysine to metabolizable energy ratio on performance and carcass characteristics of growing-finishing pigs

Pengfei Li, Zhikai Zeng, Ding Wang, Lingfeng Xue, Rongfei Zhang and Xiangshu Piao*

Abstract

A total of 2,121 growing-finishing pigs (Duroc × Landrace × Large White) were utilized in six experiments conducted to determine the effects of different ratios of standardized ileal digestible lysine (SID-Lys) to metabolizable energy (ME) on the performance and carcass characteristics of growing-finishing pigs. Exps. 1 (30 to 50 kg), 2 (52 to 70 kg) and 3 (81 to 104 kg) were conducted to find an optimum ME level and then this level was used in Exps. 4 (29 to 47 kg), 5 (54 to 76 kg) and 6 (84 to 109 kg) to test the response of pigs to different ratios of SID-Lys:ME. In Exps. 1 to 3, four treatments were used consisting of diets with a formulated ME content of 3.1, 3.2, 3.3 or 3.4 in Exps. 1 and 2 while Exp. 3 used 3.05, 3.15, 3.25 or 3.35 Mcal/kg. A constant SID-Lys:ME ratio of 2.6, 2.3 or 2.0 g/Mcal was used in Exps. 1, 2 and 3, respectively. Weight gain was significantly increased with increasing energy level in Exp. 1 while weight gain was unaltered in Exps. 2 and 3. For all three experiments, feed intake was decreased ($P < 0.05$) and feed efficiency was improved ($P < 0.05$) with increasing energy level. Tenth rib back fat thickness linearly increased ($P < 0.05$) with increasing energy level. In Exps. 4 to 6, five treatments were used consisting of diets with a SID-Lys:ME ratio of 2.4, 2.6, 2.8, 3.0 or 3.2 in Exp. 1, 2.1, 2.3, 2.5, 2.7, 2.9 or 3.2 in Exp. 2 and 1.8, 2.0, 2.2, 2.4, or 2.6 in Exp. 3. A constant ME level 3.2, 3.2 and 3.05 Mcal/kg was used in Exps. 1, 2 and 3, respectively (selected based on the results of weight gain). For all three experiments, weight gain increased ($P < 0.05$) and feed efficiency improved linearly ($P < 0.05$) as the SID-Lys:ME ratio increased. Tenth rib back fat thickness linearly decreased ($P < 0.05$) as the SID-Lys:ME ratio increased. Based on a straight broken-line model, the estimated SID-Lys:ME ratio to maximize weight gain was 3.0, 2.43 and 2.2 for 29 to 47, 54 to 76 and 84 to 109 kg of pigs, respectively.

Keywords: carcass characteristics, performance, growing-finishing pigs, metabolizable energy, standardized ileal digestible lysine

Background

Two important objectives in pig production are to maximize growth rate and improve the efficiency of nutrient utilization [1]. Dietary lysine is a key factor which influences the achievement of these objectives because it is the first limiting amino acid in diets fed to swine [2]. Dietary energy influences feed intake in growing-finishing pigs fed *ad libitum* [3,4], so that a pig's amino acid intake may be altered as the energy content of the diet changes. Therefore, it is necessary to maintain an optimum lysine to

energy ratio when the amino acid or energy content of the diet increases [1,3].

Previous studies have been conducted using variable ratios of lysine to energy [1]. Cho et al. [5] observed that the best lysine to digestible energy (DE) ratio for maximum amino acid digestibility of the pig is 2.4 g total lysine/Mcal DE for barrows (Landrace × Yorkshire × Duroc; 64 kg). Chang et al. [6] showed that the optimal lysine to DE ratios were 3.2 and 3.8 g total lysine/Mcal DE for barrows and gilts (Landrace × Yorkshire × Duroc; 16 to 57 kg), respectively. Bikker et al. [7] suggested that 2.5 g apparent ileal lysine/Mcal DE was required to optimize performance for lean gilts (20 to 45 kg). Factors which are responsible for the variation include environment and

* Correspondence: piaoxxsh@mafic.ac.cn
Ministry of Agriculture Feed Industry Centre, State Key Laboratory of Animal Nutrition, China Agricultural University, Beijing 100193, China

advance in the genetics of modern pigs [8,9]. Previous recommendation [2] is no longer fulfilling the desperate need of animal industry and production in China. Therefore, it is necessary for us to further research the lysine and energy requirement of growing-finishing pigs (Duroc × Landrace × Large White) under commercial pig farms condition in Hebei province, thereby contributing to the solid construction of Chinese feeding criterion.

Ileal digestibility coefficients for amino acids can be expressed as apparent, standardized or true. However, apparent ileal digestibility values are not always additive in a mixed diet [10]. In addition, it is difficult to measure specific endogenous amino acid losses and as a result, true ileal digestibility is not very practical for use in routine diet formulation [10]. Therefore, standardized ileal digestible values have been suggested as the best choice to be used for routine feed formulation [11] because it is additive in mixed diets and does not require the measurement of specific endogenous amino acid loss [12].

Besides the previous study on nursery pigs, the objective of this study was to determine the optimum ratio of SID-Lys:ME to maximize performance and carcass characteristics of growing-finishing pigs housed under commercial farm conditions.

Materials and methods

Animals and facilities

The experimental protocols and procedures used in these experiments were approved by the Institutional Animal Care and Use Committee of China Agricultural University (Beijing, China). The experiments were carried out at the Hebei Huailai Changfu Pig Culture Company (Hebei, China). All pigs (Duroc × Landrace × Large White) were housed in an all-in, all-out room in which the temperature was controlled between 17 and 22°C. The light schedule was 12 h light: 12 h dark. Pigs had free access to water and feed with diets provided in the form of a mash.

Experimental diets and measurements

The ME content of the corn, soybean meal and wheat bran used in these experiments (Table 1) were determined previously in our laboratory (data not published). The ME content of soybean oil (calculated as 96% of DE) was obtained from Feeding Standard of Swine [2]. The SID lysine content of corn, soybean meal and wheat bran (Table 1) were determined in our laboratory using pigs surgically equipped with a T-cannula in the distal ileum (data not published). The basal ileal endogenous losses were measured by feeding a nitrogen-free diet. The SID lysine content of L-lysine HCL lysine was assumed to be 100%. The ME and SID lysine content of the experimental diets were calculated by multiplying the ME and SID content of the individual ingredients by their inclusion level in the diets and summing the product.

Six experiments were conducted to determine the effects of different ratios of standardized ileal digestible lysine (SID-Lys) to metabolizable energy (ME) on the performance and carcass characteristics of growing-finishing pigs during the growing, early finishing and late finishing periods. Exps. 1 (30 to 50 kg), 2 (52 to 70 kg) and 3 (81 to 104 kg) were conducted to find an optimum ME level and then this level was used in Exps. 4 (29 to 47 kg), 5 (54 to 76 kg) and 6 (84 to 109 kg) to test the response of pigs to different ratios of SID-Lys:ME.

For Exps. 1 to 3, a total of 360, 312 and 264 crossbred pigs (Duroc × Landrace × Large White), weighing 30.66 ± 4.55 kg, 51.98 ± 5.45 kg and 81.08 ± 8.40 kg of BW were assigned to one of four treatments. The four treatments consisted of diets with a formulated ME content of 3.1, 3.2, 3.3 or 3.4 in Exps. 1 and 2 while Exp. 3 used 3.05, 3.15, 3.25 or 3.35 Mcal/kg. A constant SID Lys:ME ratio of 2.6, 2.3 or 2.0 g/Mcal was used in Exps. 1, 2 and 3, respectively (Table 2). Other amino acids were balanced relative to lysine using crystalline amino acids in order to match the ideal amino acid profile of Feeding Standard of Swine [2].

For Exps 4 to 6, a total of 450, 375 and 360 crossbred pigs (Duroc × Landrace × Large White), weighing 28.98 ± 5.41 kg, 53.95 ± 6.06 kg and 85.45 ± 8.14 kg of BW were assigned to one of five treatments. The treatments consisted of diets with a SID-Lys:ME ratio of 2.4, 2.6, 2.8, 3.0 or 3.2 for Exp. 1, 2.1, 2.3, 2.5, 2.7, 2.9 or 3.2 for Exp. 2 and 1.8, 2.0, 2.2, 2.4, or 2.6 for Exp. 3. A constant ME level 3.2, 3.2 and 3.05 Mcal/kg was used in Exps. 1, 2 and 3, respectively (Table 3). Other amino acids were balanced relative to lysine using crystalline amino acids in order to match the ideal amino acid profile of the Feeding Standard of Swine [2].

Each experiment used a randomized complete block design experiment which was conducted for 28 days. Each treatment was applied to six pens with 11 to 15 pigs (half male, half female) per pen. Each pen (3×4.5 m²) was equipped with a nipple waterer and two-hole dry feeder. Individual pigs and feeders were weighed at the beginning and the end of the experiment and these values were used to calculate weight gain, feed intake and feed efficiency.

At the end of Exps. 3 and 6, one pig was selected randomly from each pen to be slaughtered to determine carcass characteristics (left side of each carcass). Hot carcass weight (dressing percentage = carcass/body weight), tenth rib back fat thickness, and loin eye ($0.7 \times$ loin eye width \times depth, cm²) were measured. The pH of the carcass was measured with a pH meter (HI 8424NEW, HANNA, Rome, Italy) 45 min after slaughter and again 24 h after being placed in a 4°C refrigerator. Meat color was determined with a Chromameter, CR 400, (Minolta; Tokyo, Japan).

Table 1 ME and SID lysine values of individual ingredients used in experiments 1 to 6

	Corn	Soybean meal	Wheat bran	Soybean oil
Metabolizable energy ¹ , Mcal/kg	3.39	3.59	2.18	8.40
Standardized ileal digestible lysine ² , %	0.18	2.39	0.38	-

¹The metabolizable energy content of corn, soybean meal and wheat bran were determined previously in our laboratory (data not published). Metabolizable energy of soybean oil (calculated as 96% of DE) was obtained from Feeding Standard of Swine [2].

²Standardized ileal digestible lysine was determined previously using pigs surgically equipped with a T-cannula in the distal ileum. The basal ileal endogenous losses were measured by feeding a nitrogen-free diet (data not published).

Chemical analysis

At the beginning of each experiment, feed samples were collected and ground to pass through a 1.0-mm screen (40 mesh). Analyses for dry matter, calcium, and total phosphorus were conducted according to the methods of AOAC [13]. Gross energy was measured by an automatic adiabatic oxygen bomb calorimeter (Parr 6300 Calorimeter; Moline, IL, USA).

The amino acid concentration in diets was analyzed after the diets were ground through a 60 mesh screen. Feed samples were hydrolyzed in 6 N HCl (10 mL) at 110°C for 24 h under nitrogen. Sulphur containing amino

acids were measured after performic acid oxidation [13]. Tryptophan content was determined colorimetrically after alkaline hydrolysis following the procedures described by Miller [14]. Then the amino acids were analyzed by using a S-433D Amino Acid Analyzer (Sykam GmbH; Kleinstheim, Germany). Identification and quantification of amino acids were achieved by comparing the retention times of the peaks with those of standards.

Statistical analysis

The pen was the experimental unit for all analyses. Data were analyzed by ANOVA using a randomized complete

Table 2 Ingredient and chemical composition of experimental diets used in experiments 1 to 3 (as-fed)

Item	ME level in 30 to 50 kg, Mcal/kg				ME level in 52 to 70 kg, Mcal/kg				ME level in 81 to 104 kg, Mcal/kg			
	3.10	3.20	3.30	3.40	3.10	3.20	3.30	3.40	3.05	3.15	3.25	3.35
Ingredient, %												
Corn	57.00	63.00	70.15	68.27	58.00	65.00	72.35	71.60	60.28	67.00	75.25	77.47
Soybean meal, (43% CP)	22.50	24.00	26.00	26.00	21.00	22.08	23.00	23.00	15.50	16.50	17.00	18.00
Wheat bran	16.97	9.36	0.00	0.00	17.75	9.50	1.00	0.00	21.05	13.31	4.28	0.00
Soybean oil	0.00	0.00	0.00	1.85	0.00	0.00	0.00	1.72	0.00	0.00	0.00	0.95
Dicalcium phosphate	0.75	1.00	1.40	1.40	0.60	0.90	1.25	1.30	0.55	0.75	1.20	1.40
Limestone	1.30	1.15	0.95	0.95	1.25	1.10	0.95	0.90	1.20	1.00	0.80	0.70
Salt	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35
L-Lysine+HCl	0.13	0.14	0.15	0.18	0.05	0.07	0.10	0.13	0.07	0.09	0.12	0.13
Vitamin and mineral premix ¹	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Calculated composition												
ME, Mcal/kg	3.11	3.20	3.31	3.40	3.11	3.20	3.30	3.40	3.05	3.15	3.25	3.35
SID Lys, %	0.81	0.83	0.87	0.89	0.71	0.74	0.76	0.78	0.61	0.64	0.65	0.67
SID-Lys/ME, g/Mcal	2.60	2.60	2.60	2.60	2.30	2.30	2.30	2.30	2.00	2.00	2.00	2.00
Analyzed composition, %												
Lysine	0.92	0.94	0.97	0.99	0.83	0.84	0.86	0.88	0.72	0.73	0.74	0.76
Methionine	0.24	0.25	0.25	0.26	0.23	0.24	0.25	0.24	0.20	0.22	0.21	0.22
Threonine	0.66	0.67	0.70	0.70	0.64	0.66	0.65	0.65	0.55	0.56	0.55	0.57
Tryptophan	0.21	0.21	0.22	0.22	0.20	0.19	0.19	0.19	0.17	0.17	0.17	0.16
Calcium	0.73	0.75	0.75	0.74	0.70	0.71	0.69	0.71	0.64	0.63	0.65	0.62
Phosphorus	0.64	0.62	0.63	0.63	0.60	0.61	0.60	0.59	0.60	0.58	0.59	0.58

¹ Vitamin and mineral premix provided the following per kg of feed:

30 to 50 kg: Vitamin A, 6,000 IU; Vitamin D₃, 2,400 IU; Vitamin E, 21.6 IU; Vitamin K₃, 2 mg; Vitamin B₁, 0.96 mg; Vitamin B₂, 5.2 mg; Vitamin B₆, 2 mg; Vitamin B₁₂, 12 µg; Nicotinic acid, 22 mg; Pantothenic acid, 11.2 mg; Folic acid, 0.4 mg; Biotin, 40 µg; Choline chloride, 0.4 g; Fe, 120 mg; Cu, 140 mg; Zn, 100 mg; Mn, 16 mg; I, 0.24 mg; Se, 0.4 mg.

52 to 70 kg: Vitamin A, 5,600 IU; Vitamin D₃, 2,200 IU; Vitamin E, 21.6 IU; Vitamin K₃, 1.8 mg; Vitamin B₁, 0.88 mg; Vitamin B₂, 4 mg; Vitamin B₆, 1.8 mg; Vitamin B₁₂, 12 µg; Nicotinic acid, 20 mg; Pantothenic acid, 10 mg; Folic acid, 0.4 mg; Biotin, 40 µg; Choline chloride, 0.32 g; Fe, 88 mg; Cu, 120 mg; Zn, 96 mg; Mn, 16 mg; I, 0.24 mg; Se, 0.4 mg.

81 to 104 kg: Vitamin A, 5,200 IU; Vitamin D₃, 2,000 IU; Vitamin E, 17.2 IU; Vitamin K₃, 1.6 mg; Vitamin B₁, 0.8 mg; Vitamin B₂, 3.6 mg; Vitamin B₆, 1.6 mg; Vitamin B₁₂, 10 µg; Nicotinic acid, 17.6 mg; Pantothenic acid, 8.8 mg; Folic acid, 0.38 mg; Biotin, 32 µg; Choline chloride, 0.24 g; Fe, 76 mg; Cu, 120 mg; Zn, 76 mg; Mn, 12 mg; I, 0.24 mg; Se, 0.4 mg.

Table 3 Ingredient and chemical composition of experimental diets used in experiments 4 to 6 (as-fed)

Item	Lys/ME in 29 to 47 kg, g/Mcal					Lys/ME in 54 to 76 kg, g/Mcal					Lys/ME in 84 to 109 kg, g/Mcal				
	2.40	2.60	2.80	3.00	3.20	2.10	2.30	2.50	2.70	2.90	1.80	2.00	2.20	2.40	2.60
Ingredient, %															
Corn	63.00	63.00	63.00	63.00	63.00	65.35	65.00	65.00	65.00	65.00	60.50	60.28	60.28	60.28	60.28
Soybean meal (43% CP)	24.00	24.00	24.00	24.00	24.00	21.80	22.08	22.08	22.08	22.08	15.00	15.50	15.50	15.50	15.50
Wheat bran	9.44	9.36	9.21	9.04	8.89	9.50	9.50	9.39	9.22	9.06	21.40	21.05	20.94	20.81	20.64
Soybean oil	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Dicalcium phosphate	1.00	1.00	1.00	1.00	1.00	0.90	0.90	0.90	0.90	0.90	0.55	0.55	0.55	0.55	0.55
Limestone	1.15	1.15	1.15	1.15	1.15	1.10	1.10	1.10	1.10	1.10	1.20	1.20	1.20	1.20	1.20
Salt	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35
L-Lysine+HCl	0.06	0.14	0.22	0.30	0.38	0.00	0.07	0.15	0.23	0.32	0.00	0.07	0.15	0.22	0.30
MHA ¹	0.00	0.00	0.02	0.04	0.06	0.00	0.00	0.01	0.04	0.06	0.00	0.00	0.01	0.02	0.05
L-Threonine	0.00	0.00	0.04	0.09	0.13	0.00	0.00	0.02	0.06	0.10	0.00	0.00	0.02	0.06	0.11
L-Tryptophan	0.00	0.00	0.01	0.03	0.04	0.00	0.00	0.00	0.02	0.03	0.00	0.00	0.00	0.01	0.02
Vitamin and mineral premix ²	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Calculated composition															
SID Lys (%)	0.77	0.83	0.88	0.96	1.02	0.68	0.74	0.80	0.86	0.93	0.55	0.61	0.68	0.73	0.79
SID Lys/ME (g/Mcal)	2.4	2.6	2.8	3.0	3.2	2.1	2.3	2.5	2.7	2.9	1.8	2.0	2.2	2.4	2.6
ME (Mcal/kg)	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.19	3.05	3.06	3.06	3.05	3.05
Analyzed composition (%)															
Lysine	0.89	0.94	1.00	1.07	1.14	0.79	0.84	0.89	0.97	1.04	0.65	0.73	0.80	0.84	0.90
Methionine	0.25	0.25	0.27	0.28	0.31	0.24	0.26	0.26	0.28	0.29	0.20	0.21	0.23	0.23	0.24
Threonine	0.68	0.69	0.72	0.77	0.81	0.65	0.65	0.66	0.71	0.74	0.55	0.55	0.56	0.60	0.66
Tryptophan	0.21	0.21	0.22	0.24	0.24	0.21	0.20	0.21	0.22	0.23	0.17	0.18	0.17	0.18	0.19
Calcium	0.75	0.75	0.75	0.74	0.74	0.71	0.70	0.70	0.71	0.70	0.63	0.65	0.65	0.63	0.63
Phosphorus	0.63	0.62	0.63	0.64	0.63	0.61	0.61	0.60	0.62	0.60	0.61	0.60	0.59	0.62	0.61

¹ DL-methionine hydroxy analogue (84%), brand name of Novus International Inc., St. Louis, MO.

² Vitamin and mineral premix provided the following per kg of feed:

29 to 47 kg: Vitamin A, 6,000 IU; Vitamin D₃, 2,400 IU; Vitamin E, 21.6 IU; Vitamin K₃, 2 mg; Vitamin B₁, 0.96 mg; Vitamin B₂, 5.2 mg; Vitamin B₆, 2 mg; Vitamin B₁₂, 12 µg; Nicotinic acid, 22 mg; Pantothenic acid, 11.2 mg; Folic acid, 0.4 mg; Biotin, 40 µg; Choline chloride, 0.4 g; Fe, 120 mg; Cu, 140 mg; Zn, 100 mg; Mn, 16 mg; I, 0.24 mg; Se, 0.4 mg.

54 to 76 kg: Vitamin A, 5,600 IU; Vitamin D₃, 2,200 IU; Vitamin E, 21.6 IU; Vitamin K₃, 1.8 mg; Vitamin B₁, 0.88 mg; Vitamin B₂, 4 mg; Vitamin B₆, 1.8 mg; Vitamin B₁₂, 12 µg; Nicotinic acid, 20 mg; Pantothenic acid, 10 mg; Folic acid, 0.4 mg; Biotin, 40 µg; Choline chloride, 0.32 g; Fe, 88 mg; Cu, 120 mg; Zn, 96 mg; Mn, 16 mg; I, 0.24 mg; Se, 0.4 mg.

84 to 109 kg: Vitamin A, 5,200 IU; Vitamin D₃, 2,000 IU; Vitamin E, 17.2 IU; Vitamin K₃, 1.6 mg; Vitamin B₁, 0.8 mg; Vitamin B₂, 3.6 mg; Vitamin B₆, 1.6 mg; Vitamin B₁₂, 10 µg; Nicotinic acid, 17.6 mg; Pantothenic acid, 8.8 mg; Folic acid, 0.38 mg; Biotin, 32 µg; Choline chloride, 0.24 g; Fe, 76 mg; Cu, 120 mg; Zn, 76 mg; Mn, 12 mg; I, 0.24 mg; Se, 0.4 mg.

block design according to the GLM procedure of SAS (SAS Inst. Inc.; Cary, NC, USA). Blocks were based on initial body weight. Total variation was divided into treatment, block and error as follows:

$$Y_{ij} = \mu + T_{ij} + R_{ij} + \xi_{ij}$$

(i = 1, 2, ..., a, j = 1, 2, ..., b)

Where Y_{ij} is the observed value, μ is the population mean, T_{ij} is the treatment effect, R_{ij} is block effect, ξ_{ij} is the random error, a is the number of treatments (4 or 5), b is the number of blocks (6 in each experiment). Statistical differences among treatments were separated by Duncan's Multiple Range Test. Statistical significance was declared at $P < 0.05$ and a trend was expressed when $P < 0.10$. Results are presented as least squares

means. In Exps. 4, 5 and 6, a straight broken line regression model with a random component included for parameter L (LVAR, represents the block effect and produces different lines for the respective blocks) was used to estimate a break-point according to Robbins et al. [15] using the NLMixed procedure of SAS [16].

Results and discussion

The effects of increasing ME level on the performance of pigs during the growing, early finishing and late finishing stages are shown in Table 4. Weight gain was significantly increased ($P < 0.05$) with increasing energy level in Exp.1 (30 to 50 kg) while weight gain was unaltered in Exps. 2 (52 to 70 kg) and 3 (81 to 104 kg). But weight gain tended ($P < 0.10$) to greater in ME level of 3.2 and 3.3 than 3.1 Mcal/kg in Exp. 2. Liu et al. [17]

Table 4 Effect of dietary ME level on performance of growing-finishing pigs (Experiments 1, 2 and 3)¹

Item	ME level, Mcal/kg				SEM	P value		
	3.1	3.2	3.3	3.4		ANOVA	Linear	Quadratic
Experiment 1 (30 to 50 kg)	3.1	3.2	3.3	3.4				
Weight gain, kg/d	0.64 ^a	0.68 ^b	0.67 ^b	0.67 ^b	0.01	0.02	0.09	0.19
Feed intake, kg/d	2.10 ^a	1.92 ^{ab}	1.79 ^b	1.70 ^b	0.07	0.01	0.43	0.01
Feed efficiency	0.31 ^a	0.36 ^b	0.38 ^b	0.40 ^b	0.02	< 0.01	0.95	< 0.01
Experiment 2 (52 to 70 kg)	3.1	3.2	3.3	3.4				
Weight gain, kg/d	0.74	0.79	0.79	0.76	0.02	0.08	0.57	0.31
Feed intake, kg/d	2.75 ^a	2.57 ^{ab}	2.51 ^{ab}	2.33 ^b	0.09	0.04	0.06	0.18
Feed efficiency	0.27 ^a	0.31 ^{ab}	0.32 ^b	0.33 ^b	0.01	0.04	0.01	0.03
Experiment 3 (81 to 104 kg)	3.05	3.15	3.25	3.35				
Weight gain, kg/d	0.78	0.78	0.78	0.80	0.01	0.53	0.20	0.39
Feed intake, kg/d	3.38 ^a	3.27 ^a	3.20 ^a	2.97 ^b	0.06	< 0.01	0.02	0.06
Feed efficiency	0.23 ^a	0.24 ^a	0.25 ^a	0.27 ^b	0.01	< 0.01	< 0.01	0.03

^{a, b}Means in the same row with different superscripts are significantly different ($P < 0.05$).

¹Value represent means of six pens with 11 to 15 pigs per pen during a 28-d period.

reported increased weight gains when the dietary energy increased which is similar to the results of Du et al. [18]. Similar to the current study, De La Llata et al. [9] found pigs fed high energy levels had a significantly improved weight gain only during the growing stage and not during the finishing stage. Similarly, Beaulieu et al. [4] found that increasing the energy content of diets increased weight gain from 36 to 80 kg but pigs of heavier weight did not respond to increasing energy levels. Kerr et al. [19] observed increased weight gain when dietary net energy levels were increased from 25 to 41 kg, but not in the finishing phase (58 to 82 kg).

For all three experiments, feed intake was decreased ($P < 0.05$) and feed efficiency was improved ($P < 0.05$) with increasing energy level. This is consistent with previous reports [4,9,20] which supports the findings of the current experiment. It is well known that pigs will compensate for increases in the energy density of their diet by decreasing their feed intake [21,22].

The effects of increasing ME level on the carcass characteristics of finishing pigs are shown in Table 5. Tenth

rib back fat thickness increased linearly ($P < 0.05$) with increasing dietary ME content. Loin area showed a quadratic trend ($P = 0.06$). Loin eye area increased as the dietary ME level was increased to 3.15 to 3.25 Mcal/kg but then declined as the energy level was further increased to 3.35 Mcal/kg. Increasing the dietary ME had no effect on dressing percentage, $\text{pH}_{45 \text{ min}}$, $\text{pH}_{24 \text{ h}}$ and meat color.

In the literature, the effects of dietary energy on back fat thickness have been shown to be variable. In one experiment of the study of Beaulieu et al. [4], increasing energy did not influence back fat thickness, but in another experiment, back fat thickness increased and loin area declined as dietary energy increased, which is similar to the present study. Apple et al. [23] and Lawrence et al. [24] also observed back fat thickness increased with increasing dietary energy.

The effects of increasing SID-Lys:ME ratio on the performance of pigs during the growing, early finishing and late finishing stages are shown in Table 6. For all three experiments, weight gain increased ($P < 0.05$) and feed efficiency improved linearly ($P < 0.05$) as the SID-Lys:

Table 5 Effect of dietary ME level on carcass characteristics of finishing pigs (Experiment 3)¹

Item	ME, Mcal/kg				SEM	P value		
	3.05	3.15	3.25	3.35		ANOVA	Linear	Quadratic
Dressing percentage, %	71.98	72.31	73.30	74.03	1.86	0.86	0.34	0.64
10 th -rib back fat thickness, mm	18.5	21.5	22.3	24.0	1.42	0.09	0.02	0.07
Loin area, cm ²	39.62	44.42	42.98	40.31	1.59	0.16	0.93	0.06
$\text{pH}_{45 \text{ min}}$	6.42	6.19	6.30	6.30	0.09	0.37	0.48	0.39
$\text{pH}_{24 \text{ h}}$	5.58	5.53	5.50	5.49	0.05	0.58	0.17	0.36
L* (Lightness)	41.45	40.81	40.77	42.58	1.77	0.88	0.67	0.72
a* (Redness)	14.31	13.49	15.61	14.57	0.66	0.20	0.35	0.64
b* (Yellowness)	5.24	4.74	5.54	5.67	0.56	0.65	0.38	0.58

¹Value represent means of six pens with one pig per pen.

Table 6 Effect of dietary increasing SID-Lys:ME ratio on performance of growing-finishing pigs (Experiments 4, 5 and 6)¹

Item	SID-Lys:ME ratio, g/Mcal					SEM	P value		
	2.4	2.6	2.8	3.0	3.2		ANOVA	Linear	Quadratic
Experiment 4 (29 to 47 kg)	2.4	2.6	2.8	3.0	3.2				
Weight gain, kg/d	0.62 ^a	0.64 ^{ab}	0.64 ^{ab}	0.68 ^b	0.66 ^b	0.01	0.04	0.09	0.22
Feed intake, kg/d	1.85	1.80	1.76	1.74	1.73	0.06	0.66	0.34	0.61
Feed efficiency	0.34	0.36	0.37	0.39	0.38	0.02	0.24	0.04	0.09
Experiment 5 (54 to 76 kg)	2.1	2.3	2.5	2.7	2.9				
Weight gain, kg/d	0.73 ^a	0.77 ^b	0.80 ^b	0.80 ^b	0.79 ^b	0.01	< 0.01	0.01	0.01
Feed intake, g/d	2.44	2.44	2.38	2.35	2.34	0.08	0.86	0.49	0.79
Feed efficiency	0.30	0.32	0.34	0.34	0.34	0.01	0.16	0.05	0.10
Experiment 6 (84 to 109 kg)	1.8	2.0	2.2	2.4	2.6				
Weight gain, kg/d	0.79 ^{ab}	0.76 ^a	0.88 ^c	0.86 ^c	0.84 ^{bc}	0.03	< 0.01	0.03	0.06
Feed intake, kg/d	3.26	3.20	3.34	3.35	3.15	0.10	0.24	0.86	0.57
Feed efficiency	0.24	0.24	0.26	0.26	0.27	0.01	0.07	0.02	0.08

^{a-c}Means in the same row with different superscripts are significantly different ($P < 0.05$).

¹Value represent means of six pens with 11 to 15 pigs per pen during a 28-d period.

ME ratio increased. In the study of Smith et al. [3], increasing the lysine:ME ratio increased weight gain but did not influence feed efficiency. However, Friesen et al. [25] found weight gain increased and feed efficiency improved while feed intake tended to increase in growing pigs (34 to 72 kg).

Chiba et al. [26] and Rao and McCracken [27] both found increased weight gain and feed efficiency in growing pigs when the lysine:energy ratio increased. [1] observed increased weight gain and improved feed efficiency with increasing dietary Lys:ME ratio both in growing-finishing barrows and gilts. This is consistent with present study.

The effects of increasing SID-Lys:ME ratio on carcass characteristics of finishing pigs are shown in Table 7. Tenth rib back fat thickness was decreased (linear, $P < 0.05$; Quadratic, $P < 0.05$) as the SID-Lys:ME ratio increased. Loin area increased ($P < 0.05$) as the SID-Lys:ME ratio increased. Increasing the dietary SID-Lys/ME ratio had no effect on dressing percentage, pH_{45 min},

pH_{24 h} and meat colors. Smith et al. [3] reported that carcass characteristics were not influenced by dietary lysine to energy ratio. However, in other studies, increasing the lysine to energy ratio decreased body fat and increased loin area [1,28,29] which somewhat supports the findings of the current experiment.

The data from Exps. 4 to 6 were fitted to a straight broken-line regression equation:

$$y = 0.6664 - 0.07787 \times (3.0 - x) + \text{LVAR}$$

$$(r^2 = 0.76, \text{adjusted } r^2 = 0.64; \text{Exp. 4, Figure 1})$$

$$y = 0.7942 - 0.1933 \times (2.43 - x) + \text{LVAR}$$

$$(r^2 = 0.76, \text{adjusted } r^2 = 0.63; \text{Exp. 5, Figure 2})$$

$$y = 0.8587 - 0.209 \times (2.2 - x) + \text{LVAR}$$

$$(r^2 = 0.74, \text{adjusted } r^2 = 0.59; \text{Exp. 6, Figure 3})$$

Parameter LVAR represents the block effect and produced different lines for the respective blocks. Based on a

Table 7 Effect of dietary increasing SID-Lys:ME ratio on carcass characteristics of finishing pigs (Experiment 6)

	SID-Lys:ME ratio, g/Mcal					SEM	P value		
	1.8	2.0	2.2	2.4	2.6		ANOVA	Linear	Quadratic
Dressing percentage, %	73.04	73.25	74.25	74.99	75.33	1.62	0.81	0.18	0.42
10 th -rib back fat thickness, mm	21.00 ^a	21.67 ^a	20.00 ^a	19.67 ^a	17.17 ^b	0.82	0.01	0.02	0.03
Loin area, cm ²	40.55 ^a	40.87 ^a	41.84 ^{ab}	41.29 ^a	43.34 ^b	0.63	0.04	0.09	0.22
pH _{45 min}	6.63	6.31	6.15	6.64	6.65	0.26	0.53	0.65	0.35
pH _{24 h}	5.35	5.54	5.45	5.39	5.53	0.22	0.96	0.72	0.94
L* (Lightness)	42.81	43.09	42.01	44.5	43.05	1.71	0.89	0.73	0.94
a* (Redness)	13.48	12.97	14.7	13.35	13.89	0.75	0.56	0.58	0.80
b* (Yellowness)	5.77	5.26	6.04	6.23	6.11	0.52	0.70	0.29	0.58

^{a, b}Means in the same row with different superscripts are significantly different ($P < 0.05$).

¹Value represent means of six pens with one pig per pen.




Figure 1 Fitted straight broken-line of weight gain as a function of SID-Lys:ME ratio (Exp. 4). Observed block mean values in each treatment (dot represent block 1, square represent block 2, triangle represent block 3, x represent block 4, plus represent block 5, DIAMOND represent block 6) are shown. The straight broken line: $y = 0.6664 - 0.07787 \times (3 - x) + \text{LVAR}$ ($r^2 = 0.76$, adjusted $r^2 = 0.64$). Parameter LVAR represents the block effect and produces different lines for the respective blocks. The optimum SID-Lys:ME ratio was 3 in 28 to 47 kg pig.




Figure 2 Fitted straight broken-line of weight gain as a function of SID-Lys:ME ratio (Exp. 5). Observed block mean values in each treatment (dot represent block 1, square represent block 2, triangle represent block 3, x represent block 4, plus represent block 5, DIAMOND represent block 6) are shown. The straight broken line: $y = 0.7942 - 0.1933 \times (2.4336 - x) + \text{LVAR}$ ($r^2 = 0.76$, adjusted $r^2 = 0.63$). Parameter LVAR represents the block effect and produces different lines for the respective blocks. The optimum SID-Lys:ME ratio was 2.43 in 54 to 76 kg pig.




Figure 3 Fitted straight broken-line of weight gain as a function of SID-Lys:ME ratio (Exp. 6). Observed block mean values in each treatment (dot represent block 1, square represent block 2, triangle represent block 3, x represent block 4, plus represent block 5, DIAMOND represent block 6) are shown. Two observations (x in treatment two and DIAMOND in treatment five) were considered outliers because they were influential points in the model. The straight broken line: $y = 0.8587 - 0.209 \times (2.2 - x) + LVAR$ ($r^2 = 0.74$, adjusted $r^2 = 0.59$). Parameter LVAR represents the block effect and produces different lines for the respective blocks. The optimum SID Lys:ME ratio was 2.2 in 84 to 109 kg pig.

straight broken-line model, the estimated SID-Lys:ME ratio to optimize weight gain was 3.0, 2.43 and 2.2 for 29 to 47 kg, 54 to 76 kg and 84 to 109 kg of pigs, respectively. In addition, the estimated total Lys:ME ratio to optimize weight gain was 3.34, 2.69 and 2.61 for 29 to 47 kg, 54 to 76 kg and 84 to 109 kg of pigs, respectively.

Main et al. [1] estimated 3.14, 2.66 and 2.2 g/Mcal of total Lys:ME ratio for 43 to 70 kg, 69 to 93 kg and 102 to 120 kg barrows, respectively; and 3.23, 2.8, 2.28 and 2.2 g/Mcal of total Lys:ME ratio for 35 to 60 kg, 60 to 85 kg, 78 to 103 kg and 100 to 120 kg gilts, respectively. Yen et al. [30] estimated 3.13 and 3.23 g/Mcal of total Lys:ME ratio for 20 to 55 kg barrows and gilts respectively. Chang et al. [6] reported that the optimum total Lys:ME ratio for 16 to 57 kg barrows was 3.3 g/Mcal. Yen et al. [31] estimated 2.34 and 2.73 g/Mcal of total Lys:ME ratio for 50 to 90 kg barrows and gilts respectively. Friesen et al. [25] observed 2.78 g/Mcal of total Lys:ME ratio for 55 to 72 kg gilts. Hahn et al. [32] reported that the optimum total Lys:ME ratio for 90 to 110 kg gilts was 1.77 g/Mcal.

In conclusion, based on our results, we suggest the optimum ratio of SID-Lys:ME is 3.0, 2.43 and 2.2 g/Mcal for 29 to 47 kg, 54 to 76 kg and 84 to 109 kg of pigs, respectively.

Acknowledgements

The authors sincerely acknowledge the financial support received from the State Key Laboratory of Animal Nutrition (2004DA125184-0810) of China, National Natural Science Foundation of China (31072040), Guangdong cooperation project between Industry-Academia-Research of China (2009B090300110) and the Ministry of Science and Technology of the People's Republic of China (2006BAD12B05-10; Nyhyzx07-34). Thanks the great support of MHA from Novus international Inc., and thank Dr. Thacker P. A. (University of Saskatchewan, Canada) for excellent assistance in editing this manuscript.

Authors' contributions

PL carried out the experiments and drafted the manuscript. ZZ performed the statistical analysis. DW, LX and RZ participated in the design of the study. XP conceived of the study, and participated in its design and coordination. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 16 February 2012 Accepted: 1 March 2012

Published: 1 March 2012

References

1. Main RG, Dritz SS, Tokach MD, Goodband RD, Nelssen JL: **Determining an optimum lysine:calorie ratio for barrows and gilts in a commercial finishing facility.** *J Anim Sci* 2008, **86**:2190-2207.
2. Feeding Standard of Swine: China Agriculture Press, Beijing; 2004.
3. Smith JW, Tokach MD, O'Quinn PR, Nelssen JL, Goodband RD: **Effects of dietary energy density and lysine:calorie ratio on growth performance and carcass characteristics of growing-finishing pigs.** *J Anim Sci* 1999, **77**:3007-3015.

4. Beaulieu AD, Williams NH, Patience JF: **Response to dietary digestible energy concentration in growing pigs fed cereal grain-based diets.** *J Anim Sci* 2009, **87**(3):965-976.
5. Cho SB, Lee HJ, Chung IB, Long HF, Lim JS, Kim YY, Han IK: **Effects of dietary energy concentration and lysine on the digestible energy ratio for apparent amino acid digestibility in finishing barrows.** *Asian-Aus J Anim Sci* 2008, **21**(2):232-236.
6. Chang WH, Kim JD, Xuan ZN, Cho WT, Han IK, Chae BJ, Paik IK: **Optimal lysine:DE ratio for growing pigs of different sexes.** *Asian-Aus J Anim Sci* 2000, **13**(1):31-38.
7. Bikker P, Verstegen MW, Campbell RG, Kemp B: **Digestible lysine requirement of gilts with high genetic potential for lean gain, in relation to the level of energy intake.** *J Anim Sci* 1994, **72**(7):1744-1753.
8. Campbell RG, Taverner MR: **Genotype and sex effects on the relationship between energy intake and protein deposition.** *J Anim Sci* 1988, **66**(3):676-686.
9. De La Llata M, Dritz SS, Tokach MD, Goodband RD, Nelssen JL, Loughin TM: **Effects of dietary fat on growth performance and carcass characteristics of growing-finishing pigs reared in a commercial environment.** *J Anim Sci* 2001, **79**(10):2643-2650.
10. Stein HH, Seve B, Fuller MF, Moughan PJ, de Lange CF: **Invited review: Amino acid bioavailability and digestibility in pig feed ingredients: Terminology and application.** *J Anim Sci* 2007, **85**:172-180.
11. Stein HH, Fuller MF, Moughan PJ, Seve B, Mosenthin R, Jansman AJM, Fernandez JA, de Lange CF: **Definition of apparent, true, and standardized ileal digestibility of amino acids in pigs.** *Livest Sci* 2007, **109**:282-285.
12. Stein HH, Pedersen C, Wirt AR, Bohlke RA: **Additivity of values for apparent and standardized ileal digestibility of amino acids in mixed diets fed to growing pigs.** *J Anim Sci* 2005, **83**:2387-2395.
13. AOAC: **Official methods of analysis.** *Association of Official Analytical Chemists.* 16 edition. Washington, DC; 1990.
14. Miller EL: **Determination of the tryptophan content of feedingstuffs with particular reference to cereals.** *J Sci Food Agric* 1967, **18**:381-386.
15. Robbins KR, Saxton AM, Southern LL: **Estimation of nutrient requirements using broken-line regression analysis.** *J Anim Sci* 2006, **84**(Suppl):E155-165.
16. SAS: **SAS User's Guide: Statistics (Version 8.01).** SAS Inst. Inc., Cary, NC; 1999.
17. Liu ZH, Yang FY, Kong LJ, Lai CH, Piao XS, Gu YH, Ou XQ: **Effects of dietary energy density on growth, carcass quality and mRNA expression of fatty acid synthase and hormone-sensitive lipase in finishing pigs.** *Asian-Aus J Anim Sci* 2007, **20**:1578-1593.
18. Du W, Li YJ, Zhao GY, Yin YL, Kong XF: **Effect of dietary energy level on nutrient utilization, insulin-like growth factor-i and insulin-like growth factor binding protein-3 in plasma, liver and longissimus dorsi muscle in growing-finishing pigs using soybean oil as an energy source.** *Asian-Aus J Anim Sci* 2009, **22**:1180-1185.
19. Kerr BJ, Southern LL, Bidner TD, Friesen KG, Easter RA: **Influence of dietary protein level, amino acid supplementation, and dietary energy levels on growing-finishing pig performance and carcass composition.** *J Anim Sci* 2003, **81**:3075-3087.
20. Myer RO, Lamkey JW, Walker WR, Brendemuhl JH, Combs GE: **Performance and carcass characteristics of swine when fed diets containing canola oil and added copper to alter the unsaturated:saturated ratio of pork fat.** *J Anim Sci* 1992, **70**:1417-1423.
21. Cole DJA, Chadd SA: **Voluntary food intake of growing pigs.** In *The Voluntary Food Intake of Pigs.* Edited by: Forbes JM, Varley MA, Lawrence TLJ. British Society of Animal Production; 1989:61-70.
22. NRC: **Nutrient Requirements of Swine.** Natl Acad Press, Washington, DC; 10 1998.
23. Apple JK, Maxwell CV, Brown DC, Friesen KG, Musser RE, Johnson ZB, Armstrong TA: **Effects of dietary lysine and energy density on performance and carcass characteristics of finishing pigs fed ractopamine.** *J Anim Sci* 2004, **82**(11):3277-3287.
24. Lawrence BV, Adeola O, Cline TR: **Nitrogen utilization and lean growth performance of 20 to 50 kilogram pigs fed diets balanced for lysine: energy ratio.** *J Anim Sci* 1994, **72**:2887-2895.
25. Friesen KG, Nelssen JL, Goodband RD, Tokach MD, Unruh JA, Kropf DH, Kerr BJ: **Influence of dietary lysine on growth and carcass composition of high-lean-growth gilts fed from 34 to 72 kilograms.** *J Anim Sci* 1994, **72**:1761-1770.
26. Chiba LI, Lewis AJ, Peo ER Jr: **Amino acid and energy interrelationships in pigs weighing 20 to 50 kilograms: I. Rate and efficiency of weight gain.** *J Anim Sci* 1991, **69**(2):694-707.
27. Rao DS, McCracken KJ: **Effect of protein intake on energy and nitrogen balance and chemical composition of gain in growing boars of high genetic potential.** *Anim Prod* 1990, **51**:389-397.
28. Castell AG, Cliplef RL, Poste-Flynn LM, Butler G: **Performance, carcass and pork characteristics of castrates and gilts self-fed diets differing in protein content and lysine:energy ratio.** *Can J Anim Sci* 1994, **74**:519-528.
29. Cameron ND, Penman JC, Fisker AC, Nute GR, Perry AM, Wood JD: **Genotype with nutrition interactions for carcass composition and meat quality in pig genotypes selected for components of efficient lean growth rate.** *Anim Sci* 1999, **69**(1):69-80.
30. Yen HT, Cole DJA, Lewis D: **The response of pigs from 25 to 55 kg live weight to dietary ideal protein.** *Anim Prod* 1986, **43**:141-154.
31. Yen HT, Cole DJA, Lewis D: **The response of pigs from 50 to 90 kg live weight to dietary ideal protein.** *Anim Prod* 1986, **43**:155-165.
32. Hahn JD, Biehl RR, Baker DH: **Ideal digestible lysine level for early- and late-finishing swine.** *J Anim Sci* 1995, **73**:773-784.

doi:10.1186/2049-1891-3-9

Cite this article as: Li et al.: Effects of the standardized ileal digestible lysine to metabolizable energy ratio on performance and carcass characteristics of growing-finishing pigs. *Journal of Animal Science and Biotechnology* 2012 **3**:9.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

