

CORRECTION

Open Access



# Correction to: Residual feed intake phenotype and gender affect the expression of key genes of the lipogenesis pathway in subcutaneous adipose tissue of beef cattle

Clare McKenna<sup>1,2</sup>, Richard K. Porter<sup>2</sup>, Kate A. Keogh<sup>1</sup>, Sinead M. Waters<sup>1</sup>, Mark McGee<sup>1</sup> and David A. Kenny<sup>1\*</sup>

## Correction

In the original publication of this article [1], some errors in Table 4 need to be corrected as below:

“Creating, umil/L” should be “Creatine, umol/L”

“Glucose, nmlo/L” should be “Glucose, nmol/L”

“Urea, nmo/L” should be “Urea, nmol/L”

The original publication has been corrected.

## Author details

<sup>1</sup>Animal and Bioscience Research Department, Teagasc Grange, Dunsany, Meath C15 PW93, Ireland. <sup>2</sup>School of Biochemistry & Immunology, Trinity College Dublin, Dublin 2 D02 R590, Ireland.

Received: 19 October 2018 Accepted: 22 October 2018

Published online: 07 November 2018

## Reference

1. McKenna, et al. Residual feed intake phenotype and gender affect the expression of key genes of the lipogenesis pathway in subcutaneous adipose tissue of beef cattle. *J Anim Sci Biotechnol.* 2018;9:68 <https://doi.org/10.1186/s40104-018-0282-9>.

\* Correspondence: [david.kenny@teagasc.ie](mailto:david.kenny@teagasc.ie)

<sup>1</sup>Animal and Bioscience Research Department, Teagasc Grange, Dunsany, Meath C15 PW93, Ireland

Full list of author information is available at the end of the article

